

Undecidability in \mathbf{R}^n : Riddled Basins, the KAM Tori, and the Stability of the Solar System*

Matthew W. Parker^{†‡}

Some have suggested that certain classical physical systems have undecidable long-term behavior, without specifying an appropriate notion of decidability over the reals. We introduce such a notion, *decidability in μ* (or *d- μ*) for any measure μ , which is particularly appropriate for physics and in some ways more intuitive than Ko's (1991) *recursive approximability* (r.a.). For Lebesgue measure λ , *d- λ* implies r.a. Sets with positive λ -measure that are sufficiently "riddled" with holes are never *d- λ* but are often r.a. This explicates Sommerer and Ott's (1996) claim of uncomputable behavior in a system with riddled basins of attraction. Furthermore, it clarifies speculations that the stability of the solar system (and similar systems) may be undecidable, for the invariant tori established by KAM theory form sets that are not *d- λ* .

1. Introduction. Several authors have suggested that the long-term behavior of some deterministic physical systems, or of some classical models in \mathbf{R}^n , is in some sense uncomputable (Moore 1990, 1991; Moser 1978, 67–68; Pitowsky 1996; Sommerer and Ott 1996; Wolfram 1985). At their most explicit, these authors argue that the set of real-valued states leading eventually to a certain kind of behavior is undecidable. However, none of these

*Received September 2002; revised December 2002.

†This paper won the PSA 2002 Graduate Student Essay Contest. To contact the author, please write to 1719 N. Mariposa Ave., Los Angeles, CA 90027 (or to Department of Philosophy, University of Chicago, 1010 E. 59th Street, Chicago, IL 60637); e-mail: mwparker@uchicago.edu.

‡I thank Wayne Myrvold, Howard Stein, William Wimsatt, Greg Lavers, Matthew Frank, Eric Schliesser, James Meiss, Norman Leibowitz, John Sommerer, Cristopher Moore, Jim Guszcza, Jill North, Amit Hagar, Joan Wackerman, David Malament, and the referees for comments on this and related efforts, the University of Western Ontario for the opportunity to present this research, Waid Parker for financial support, and Laurel Parker.

Philosophy of Science, 70 (April 2003) pp. 359–382. 0031-8248/2003/7002-0008\$10.00
Copyright 2003 by the Philosophy of Science Association. All rights reserved.

authors gives a rigorous definition of decidability for sets of real-valued states, and this warrants more concern than one might expect.

Intuitively, a set is decidable if there is a mechanical procedure that will always determine whether or not a given object in the domain of discourse is in that set. For sets of natural numbers, this intuitive notion seems to be captured by the rigorous mathematical concept of a “recursive” set, and this concept is nontrivial in extension. That is, there are many recursive sets of natural numbers and many nonrecursive ones (setting aside any constructivistic denial of nonrecursive sets).

For sets of real numbers or real n -tuples, however, there is no uniquely standard notion of a decidable set. There *is* a standard notion of a computable *function* on the reals or real n -tuples, called *Grzegorzcyk-computability*, with many different formulations (Grzegorzcyk 1955a, b, 1957; Ko 1991; Pour-El and Caldwell 1975; Pour-El and Richards 1983; Weihrauch 2000), but it does not directly suggest a useful concept of a decidable *set* of reals. One might like to say that a set B of reals is decidable if the characteristic function of B (i.e., $\chi_B(x) = 1$ if $x \in B$, $\chi_B(x) = 0$ otherwise) is Grzegorzcyk-computable. However, this notion is practically unsatisfiable. Grzegorzcyk-computable functions are always continuous (Grzegorzcyk 1955a), but in \mathbf{R}^n , only \emptyset and \mathbf{R}^n have continuous characteristic functions.¹ Hence in the obvious sense for sets of real n -tuples, undecidability is trivial.

Myrvold (1997) provides reasons to accept this concept of decidability on \mathbf{R}^n , uninformative as it may be, and concede that nontrivial sets of reals are simply not decidable. Nonetheless, some sets of reals are more nearly decidable than others, and various relaxed notions of a decidable set of reals have been introduced (e.g., Blum, Shub, and Smale 1989; Ko 1991; Myrvold 1997; Weihrauch 2000). Most of them are far from equivalent.

Here we discuss Ko’s (1991) notion of a *recursively approximable* (r.a.) set, and a previously overlooked notion, which we dub *decidability in μ* (abbreviated $d\text{-}\mu$), where μ is a measure. Both notions involve measures and express the desire for a decision algorithm with a high probability of success. Ko’s notion has intuitive and practical appeal, and we will make some use of it here, but it also has certain features for which the motivations are unclear. We will see that $d\text{-}\mu$ is closer to the classical concept of a decidable set of natural numbers and that it captures certain intuitions that r.a. does not. In particular, r.a. implies the existence of a decision procedure that succeeds with probability *arbitrarily close* to one (given certain assumptions about the relevant probability measure), while $d\text{-}\mu$ expresses the existence of a decision procedure that succeeds with a probability of *exactly* one.

1. One can extend Grzegorzcyk-computability to functions on other metric spaces (Myrvold 1997), and in disconnected spaces there *are* nontrivial sets with continuous, computable characteristic functions.

Applications of $d-\mu$ are suggested by some interesting results in dynamical systems. A *dynamical system* is a mathematical model consisting of a space of states—*phase space*—and a flow function, by which the state of the system at any time determines the state at any other time. Phase space enables one to represent the values of all the variables of a system—such as the position and momentum coordinates for several bodies—as a single point. The *basin* of a set A of states is the set of states from which the system will asymptotically approach A . A basin is *riddled*, as in ‘riddled with holes,’ if every open set in phase space contains a positive-measure portion of the complement of the basin (Alexander et al. 1992).

Classical models of physical systems have shown numerical evidence of riddled basins (Sommerer and Ott 1993, 1996) and some actual systems have shown observational evidence of riddling or approximate² riddling (Heagy, Carroll, and Pecora 1994). A riddled basin implies a kind of unpredictability, since *exact* initial data are required in order to determine whether the state of a system lies in such a basin, and hence to determine the system’s qualitative behavior as time increases without bound. (Note this is different from “chaos,” where *very precise* initial data are required to determine *finite-time* behavior.) What is more, any *computation* that determines the long-term behavior of a system with riddled basins must *use* the complete exact initial data, which generally cannot be finitely expressed. Hence such computations are intuitively impossible, even if the data are somehow available. On this basis, Sommerer and Ott (1996) argue that a certain system that seems to have riddled basins exhibits “uncomputable” behavior.

However, the authors do not give a definition of ‘uncomputable set’ sufficient to distinguish it from the trivially satisfied notion mentioned above. Here we clarify and bolster their claim with a simple theorem: *no riddled set with positive λ -measure is $d-\lambda$* (Theorem III). (Throughout this paper, λ denotes Lebesgue measure, the standard notion of volume in \mathbf{R}^n .) Therefore if Sommerer and Ott are correct that their basins are riddled and have positive measure, the basins are undecidable in the precise sense to be defined here. Hence Sommerer and Ott’s uncomputability claims are warranted, and since the proof of this result and the motivations for $d-\mu$ are similar to Sommerer and Ott’s arguments and motivations, it seems that decidability in μ captures the intuitions of at least two scientists.

Another application is the famous problem of the stability of the solar system. There one models the solar system as point masses under Newtonian gravitation, and the problem is to determine whether any of the

2. The evidence here supports a classical model with riddled basins, but of course such models break down at the quantum level. Hence the evidence only supports approximate riddling, or riddling in the classical limit.

bodies will ever escape the system or collide with another body. If not, we say that the orbit of the whole system in phase space is *stable*. (Though, there are other notions of stability that it might not satisfy.) Such exalted mathematicians as Lagrange, Laplace, and Poisson have obtained partial stability results (see Poincaré 1891, 1898; Moser 1978; Diacu and Holmes 1996 for nontechnical reviews), but Poincaré famously revealed formidable obstacles to any complete solution: the nonexistence of any constants of motion beyond the known few, and the terribly complex interweaving of orbits now known as the homoclinic tangle (1890, [1892–1899] 1993; see also Moser 1978; Goroff’s introduction to Poincaré [1892–1899] 1993; Diacu and Holmes 1996; Parker 1998). Some have suggested that the stability problem may even be undecidable (Moser 1978; Wolfram 1985), but again, without articulating an appropriate nontrivial sense.

The concept of decidability in μ fills this gap. In fact, modern KAM theory (named for Kolmogorov, Arnol’d, and Moser), provides reasons to suspect that the stability problem and many related problems are not $d\text{-}\lambda$ (where again, λ is Lebesgue measure).³ It shows that certain classes of energy-conserving systems have many bounded orbits, confined to tori in phase space. We will see that the tori of bounded orbits established by KAM theory for a given system form a set that is not $d\text{-}\lambda$. This itself is a meaningful undecidability result for an important field, but moreover it suggests that the stability of the solar system (and similar problems) may not be $d\text{-}\lambda$. The latter proposition depends on yet unknown facts (see Section 6), but in any case our discussion will shed some light on the *senses* in which such problems might be rigorously unsolvable.

It should be said, in fairness, that the authors cited above (Moore, Pitowsky, Moser, Sommerer and Ott, and Wolfram) never set out to define a rigorous and viable concept of decidability for sets in real space. However, the issue is not whether the machines to which they refer provide a basis for such a definition (as oracle Turing machines will for us). These authors have made claims and suggestions of *undecidable* behavior for real-valued models of physical systems—they suggest not just that, for example, none of Moore’s machines can decide the fate of certain Moore machines, but that *nothing* within the traditional conception of a computing machine or algorithm can decide the fate of certain Moore machines, or of other real-valued systems. Since taken in the obvious way this is trivial, such suggestions are not clear until relevant nontrivial no-

3. Though Lebesgue measure is not always well-defined on the phase space of a mechanical system, planetary systems have natural phase spaces in \mathbf{R}^n , and the classic KAM theorems (e.g., Arnol’d 1963a, b, Moser 1973) assume contexts in which Lebesgue measure is well-defined. It also seems likely that our results will extend to other spaces and measures.

tions of undecidability have been identified. Our goal here is not to indict others for making vague claims, but to give *sharper teeth* to undecidability claims by clarifying what precisely they could mean.⁴ Technical definitions and proofs appear in the appendix.

2. Defining Decidability. We adopt the usual discrete conception of computation as the systematic manipulation of finite symbol strings. Though analog computation is worth studying, we restrict our attention to the discrete approach, which dominates mathematical practice and has been canonized by recursion theory.

In order to regard points in some space⁵ as subjects of discrete computation, we must represent them using finite strings. There being no way to code each point in an uncountable space with a single finite string, we represent real points by approximation. Fix a finite alphabet A and a coding of the natural numbers, i.e., a map taking the finite strings from A onto \mathbf{N} . Similarly, fix a coding of the rational n -tuples.⁶ We represent a point $x \in \mathbf{R}^n$ by a function from strings to strings that converges quickly to x in the following sense: it takes a code for any natural number m to a code for a rational n -tuple q such that $\|q - x\| < 2^{-m}$. Let us call such a function a *Cauchy oracle* for x (Definition 1(i) in Appendix). Note that a given point is represented by many different Cauchy oracles, for there are uncountably many sequences of rational points that converge quickly to a given point.⁷

4. Some conclude (in private communications) that Moore's (1990, 1991) arguments show certain naturally interesting sets are not decidable in λ , or not decidable disregarding boundaries in the sense of (Myrvold 1997). However, I fear there are subtle complications involved in such inferences—and in Moore's constructions—that would be better addressed in another paper.

5. We consider computation only on \mathbf{R}^n , but the concepts here easily extend to separable metric spaces, including domains important to physics such as curved manifolds and Hilbert spaces.

6. Our undecidability results are independent of these codings, but positive decidability results require somewhat informative codings. Theorems II and IV assume that the relation $\|Q_n(s_1) - Q_n(s_2)\| \leq Q_n(s_3)$ is recursive.

7. It may seem strange to represent real points in this way rather than using, say, the usual base- n digital expansions. We do so for the sake of comparability with Ko (1991). In fact, base- n expansions would be sufficient for our study of decidability concepts, but in Ko's (1991) and Weirauch's (2000) broader studies of computable *functions* and computational *complexity* on the reals, base- n expansions lead to counterintuitive results: simple operations like addition turn out to be noncomputable. In adopting Cauchy oracles as our standard representations of real n -tuples, we follow Ko except in one detail: Ko defines a "Cauchy function" with inclusive inequality (" \leq ") in place of our strict inequality (" $<$ "). The strict inequality gives Theorem I a more elegant form (see note 8), and it does not change the extension of either r.a. or d- μ .

For applications to empirical science, a Cauchy oracle represents an infinite sequence of ever more accurate measurements of a particular quantity. It embodies the idealizing assumption that with sufficient time, effort, and funding, scientists can exact a measurement to any desired accuracy, short of perfect. Unrealistic as it is, this idealization enables us to abstract away from the specific limitations of various measurement techniques, and to show what cannot be computed *no matter how accurate* the data.

We use the term ‘oracle’ here because our decidabilities will be defined in terms of *oracle Turing machines*. A *Turing machine* is a hypothetical device consisting of an infinite tape (the *work tape*) and a head that moves back and forth along the tape reading and writing symbols (say from our alphabet A). The head has finitely many possible internal states, and at each step, its motion, the symbol it writes (if any), and the next internal state it takes on are determined by (i) its present internal state, (ii) the symbol immediately under the head, and (iii) its program (a finite list of instructions that determines its response to each symbol in each state). This model can be treated rigorously, and it is well known that the functions it can compute on \mathbf{N} are precisely the *recursive* and *partial recursive* functions—the same functions picked out by several other notions of computability (Turing 1936; and, e.g., Davis and Weyuker 1983; Soare 1987). This supports the *Church-Turing thesis* that such functions are precisely the intuitively computable functions on \mathbf{N} .

An *oracle Turing machine* (OTM) is a Turing machine that can “ask questions” of a hypothetical black box called an *oracle*. In particular, a *function OTM* is a Turing machine with a second tape (the *query tape*) where in the course of its computations it may write a query in the form of a finite string, and the oracle will then replace that string with another finite string (Figure 1). Hence the oracle is just a function from finite strings to finite strings. While it plays a role *in* a machine, the oracle itself is not a machine and it need not be a computable function. In classical recursion theory, an uncomputable oracle increases an OTM’s computing power, and given a particular oracle, one considers what functions on \mathbf{N} become computable (Soare 1987).

Here, however, we do not use an oracle as an aid to computation but as an input, a representation of the real-valued point on which we wish to perform a computation. When we consider whether an OTM M correctly performs some computation on a point x , we assume that M ’s oracle is a Cauchy oracle for x . M can ask this oracle for an approximation of x with any specified accuracy, and the oracle will provide it. We then ask, “If M is provided with such an oracle, will it give an output appropriate to x ?” “If M is provided a Cauchy oracle for y , will it give an output appropriate to y ?” etc. Thus the oracle defines the particular instance of a problem that M must solve. The oracle may still be an uncomputable string func-

An oracle Turing machine M




Figure 1. A function oracle Turing machine. When the machine M enters the query state q_i , the finite string on the query tape is replaced by another, as determined by the oracle function ϕ ; the read/write head returns to the “origin,” where the new string begins; and M enters the answer state q_j . If ϕ is a Cauchy oracle for x and the string on the query tape is a code for, say, 11, then the new string codes a rational point with a distance from x less than 2^{-11} .

tion; after all, for us the oracle represents a sequence of ever more accurate measurements of some real-world quantity, and there is no reason to expect such a sequence to be computable. Neither, though, is there any worry that an uncomputable oracle will make an OTM inappropriately powerful, for we *treat* oracles as inputs: given a particular function *on* oracles, we ask *on which* oracles does an OTM compute that function.

We omit further details of OTMs and adopt the standard assumption that any systematic procedure involving queries to an oracle can be carried out by an OTM. This is the *relativized Church-Turing thesis* used in higher recursion theory (Soare 1987). It licenses us to prove computability results by appeal to informal algorithms rather than complicated Turing programs.

Both Ko’s (1991) recursive approximability and our decidability in μ are defined in terms of OTMs and measures. Ko’s notion amounts to the existence of an approximately accurate decision algorithm in the form of

an OTM such that the set of points where the OTM errs (the *error set*) can be made arbitrarily small in Lebesgue measure (or outer measure). The upper bound for the measure of the error set can be specified as an input on the machine's work tape. Also, Ko requires this machine to halt (finish computing and give an output) on every Cauchy oracle. In sum, a set is *recursively approximable* (r.a.) if there is an OTM that (1) halts on every Cauchy oracle and (2) computes the set's characteristic function correctly, except perhaps on some set with Lebesgue outer measure less than 2^{-n} for specified n (Definition 3 in Appendix).

Ko explains that such a machine would decide a set "with an error probability less than or equal to 2^{-n} , where the probability is measured by the natural Lebesgue measure" (1991). However, this motivating remark raises a few questions.

Firstly, is it legitimate to equate probability with Lebesgue measure? The connection between probability and Lebesgue measure, or the closely related "microcanonical" measure on an energy surface, has been much discussed by philosophers in relation to thermodynamics and statistical mechanics (Poincaré 1907; Sklar 1973, 1993; Malament and Zabell 1980; Batterman 1998; Vranas 1998) and is still debated. Yet, that discussion at least illustrates how attractive it is to relate probabilities to Lebesgue measure or something much like it—in particular to suppose that, barring special circumstances, the probability associated with a measure-zero set of states in phase space is zero. We adopt this assumption here as a motivation both for r.a. and for our results below involving $d-\lambda$, where λ is Lebesgue measure. However, $d-\mu$ will be defined for *any* measure μ , and even r.a. generalizes easily to an arbitrary measure or outer measure. Hence these decidability concepts, if not the results below, can be applied to the appropriate probability measure whatever it may be.

Secondly, why require a machine that always halts? Assuming we have a machine that sometimes gives incorrect output, the epistemological situation would seem no worse if in principle that machine could also fail to halt, but with probability zero. This would not affect the probability of obtaining a correct output, and in application we could be confident that nonhalting cases would never arise.

Finally, why should one be satisfied with an arbitrarily small, possibly nonzero probability of error? A probability of error exactly equal to zero would be intuitively better.

In short, why not simply demand a machine that, with probability one (whatever the probability measure may be), will halt and give correct output? With this in mind, we say a set is *decidable in μ* (or $d-\mu$), where μ is a measure, if some OTM will compute its characteristic function except perhaps on some set with μ -measure zero, where the OTM might decide incorrectly or not at all (Definition 4 in Appendix). Otherwise, the set is

undecidable in μ (or $u\text{-}\mu$). We will see below that r.a. and $d\text{-}\mu$ are not equivalent, and in particular $d\text{-}\lambda$ is strictly stronger than r.a.

Our intent in defining $d\text{-}\mu$ is that μ should be chosen to reflect the likelihood of a physical system taking on a state in a given set, at least to some extent. Specifically, μ should be a measure in which the relevant probability is *absolutely continuous*—assigning probability zero to all measure-zero sets. Then if a given set is $d\text{-}\mu$, there is a machine M such that the probability of a state arising that M does not correctly decide is zero. Though it might still be strictly *possible* for such a state to occur, we can safely assume that none will (and this assumption is reasonable whether the probability in question is an objective fact or merely a reflection of our expectations).

It is not claimed here that $d\text{-}\mu$ is the best concept of decidability in every respect for every real-valued context. In fact, r.a. is in some ways more pragmatic, for an extremely minuscule probability of error is usually good enough for practical purposes. However, it is just this pragmatism that makes r.a. less analogous to classical decidability than $d\text{-}\mu$, for the classical concept of decidability in discrete recursion theory (or for that matter, logic [Gödel 1931]) is highly theoretical. It concerns what can be decided in *all* cases using a single algorithm (or effectively axiomatized theory). Though only trivial sets of reals are decidable in this absolute sense, clearly those sets that are decidable all the way up to measure zero come closer to that standard than those that can only be decided up to an arbitrarily small nonzero measure.

One could define still stronger notions of decidability, which might be preferable in some respects. One strengthening is obtained by appending to the definition of $d\text{-}\mu$ the requirement that M must halt on every Cauchy function. However, this is too strong; if μ is a reasonably nice measure, only measure-zero sets and their complements are decidable in this sense (Proposition 2). Hopefully other strengthenings will be explored in future writings. Strictly speaking though, no stronger decidability could entail any greater *probability* of correct output. If we can indeed safely assume that barring special circumstances, no state in a given measure-zero set will actually arise, then $d\text{-}\mu$ already guarantees us an algorithm that in practice will always succeed. Also, a stronger notion of decidability would imply a weaker notion of *undecidability*, and this would be undesirable, for one of our tasks here is to find the most meaningful *undecidability* results possible for certain dynamical systems.

3. The Topological Use Principle. All of the results here are based on the Use Principle of classical recursion theory. This states that if an OTM halts, then it does so after finitely many steps and after scanning only finitely many symbols provided by the oracle (see, e.g., Soare 1987). In

our context this implies that an OTM must make its “decisions” based on approximate input, without ever “knowing” the exact position of a point. Consequently, if M outputs q on a given point, then it outputs q on a neighborhood of that point. More precisely,

Theorem I (Topological Use Principle). If an OTM M halts with output q on some Cauchy oracle for x , then there is a neighborhood U of x such that M halts with output q on some Cauchy oracle for each point in U .⁸ (See Appendix for proof.)

This implies that if a set B is d - μ as witnessed by an OTM M , and M outputs, say, 1 (for “Yes, $x \in B$ ”) on some Cauchy oracle for x , then we can determine a neighborhood around x that is contained in B except perhaps for a subset of measure zero. If we have chosen a reasonable coding of rationals, this enables us to show

Theorem II. D - λ implies r.a.

For, given an algorithm M that shows B is d - λ , we can effectively construct two sequences of neighborhoods: some on which M halts and which nearly fill \mathbf{R}^n , and others covering the gaps. It is then simple to give an algorithm that halts on all Cauchy oracles and correctly decides B almost everywhere except on the latter neighborhoods, which form an arbitrarily small set.

4. Riddled Sets. In some work on dynamical systems, a set B is called *riddled* (Alexander et al. 1992) if it is significantly pervaded with holes, i.e., if its complement has positive Lebesgue measure (λ) in every nonempty neighborhood (Definition 5; see Figure 2). For example, any nowhere-dense subset of \mathbf{R}^n is riddled, but one can also construct dense riddled sets.


It follows from the Topological Use Principle that no algorithm correctly computes the characteristic function of a set $B \subseteq \mathbf{R}^n$ at the boundary of B . Of course, a riddled set is entirely included in its own boundary. This makes it possible to show:

Theorem III. Every riddled set with positive Lebesgue measure is u - λ . (See Figure 2.)

As we will see, this explicates and justifies Sommerer and Ott’s uncomputability claims, and it also shows that the KAM tori are u - λ .

Consider a simple example of a riddled set, a *generalized Cantor set*. This is constructed in stages: beginning with the unit interval, remove an interval from the middle, then remove a much smaller interval from each remaining piece, and repeat (Figure 3). If the intervals removed decrease

8. If “ $<$ ” in the definition of a Cauchy oracle is replaced with “ \leq ” as in Ko (1991), then “of x ” in Theorem I must be replaced with “whose closure contains x .”



If M decides that $x \in A$ then M errs here.

Figure 2. Riddling and Theorem III. Here A is riddled: every neighborhood, like the magnified inset, contains positive-measure portions of the complement A^c . If an OTM M gives output q given some Cauchy oracle for x , it will also do so for each point in a small neighborhood of x (Theorem I). Therefore it incorrectly decides A on a positive-measure set within that neighborhood (Theorem III).




Figure 3. Construction of a generalized Cantor set C with positive Lebesgue measure.

rapidly in size, the set remaining after infinitely many stages will have positive measure. Clearly such a set is riddled, and if the construction is sufficiently systematic, the resulting set is r.a. Hence, in virtue of Theorem III, we obtain the following:

Theorem IV. There exists a set that is r.a. but not d - λ .

With Theorem II, this shows that d - λ is strictly stronger than r.a., and the generalized Cantor set serves as a simple model for the kind of undecidability that appears to occur in Sommerer and Ott's example and the n -body stability problem.

5. Riddled Basins of Attraction. Sommerer and Ott's model (1996) consists of a point particle in a two-dimensional potential, with an additional force given as a sinusoidal function of time. The motion is governed by

$$d^2(x, y)/dt^2 = -\gamma d(x, y)/dt - \nabla V(x, y) + \mathbf{i} a \sin(\omega t), \quad (1)$$

where γ is the friction coefficient, \mathbf{i} is the unit vector in the positive x direction, a is the amplitude of the periodic force $\mathbf{i} a \sin(\omega t)$, $\omega/2\pi$ gives the frequency of the periodic force, and $\nabla V(x, y)$ is the gradient of the potential given by

$$V(x, y) = (1 - x^2)^2 + sy^2(x^2 - p) + ky^4. \quad (1a)$$

(See Figure 4.) The parameters s , p , and k may be varied to obtain a family of potentials. With fixed parameters, the solutions of (1) form a dynamical system on a five-dimensional phase space: two dimensions to represent the position of the particle, two for momentum or velocity, and since the periodic force depends on time, we include time itself as a state variable.

Sommerer and Ott refer to Milnor's (1985) definition of *attractor*—essentially, a set whose basin of attraction has positive Lebesgue measure.⁹ They give an analytic argument that their system (with chosen parameters) has at least two attractors, one in each well of the potential. Numerically approximated graphs seem to show the disjoint basins of both attractors occupying significant portions of each neighborhood in phase space, suggesting that the basins are riddled (Figure 5). The authors also remark that the full measure of the phase space is divided between these two basins,¹⁰ implying that the basins have positive measure. Inferring that a

9. Under other definitions of attractor, all points near an attractor lie in its basin. So defined, an attractor cannot have a riddled basin. Sommerer and Ott actually define attractor differently from Milnor, but immaterially so.

10. This is plausible, but not obvious. The shape of the potential V shows that there is no "attractor at infinity," but there could be a positive-measure set of orbits that do not approach either attractor.




Figure 4. Sommerer and Ott's potential. Motion defined by equation (1) can be visualized as a marble rolling on this surface while the surface rocks to the left and right, representing the periodic force $\mathbf{ia} \sin(\omega t)$.

computation must make full use of *exact* data in order to determine membership in one of these basins, Sommerer and Ott conclude that the basins are uncomputable.

Theorem III implies that if indeed Sommerer and Ott's basins are riddled and have positive measure, then neither basin is d - λ . This seems to reflect the intuitions behind Sommerer and Ott's uncomputability claim, for what makes the undecidability of their example strong is the fact that the complement of a riddled set has *positive measure* in every neighborhood. Therefore any algorithm will fail to decide it not just in a few isolated cases, but on a set of cases with positive measure and, intuitively, with a positive probability that some case in that set will actually occur. This is precisely the worry that motivated our definitions of d - μ and u - μ . Also, Sommerer and Ott's intuition that an algorithm cannot actually *use* infinitely precise data about the position of a point is essentially the Topological Use Principle, the main insight used to establish Theorem III. Hence u - λ seems to be roughly the undecidability they had in mind. Regardless, it is one rigorously defined undecidability that follows from their claims.




Figure 5. Sommerer and Ott's riddled basins (1996). Slices of the five-dimensional phase space of the dynamical system defined by equation (1). The attractors intersect these planes along the x -axis. For each initial condition in each 760×760 grid, Sommerer and Ott simulated an orbit until it came within 10^{-5} of an attractor, with velocity transverse to the attractor less than 10^{-6} . Initial states leading to the left attractor were colored black, and those leading to the right attractor, white. Blow-ups (b) and (c) of insets in (a), and (d) of the inset in (c), suggest that both basins are riddled.

On the other hand, r.a. is apparently not the concept of computability Sommerer and Ott had in mind when claiming that their basins were uncomputable. Rather, r.a. explicates their claims that the basins are in a sense computable—sufficiently computable that the authors could produce qualitatively accurate graphs revealing the riddled structure of those basins. In fact, Sommerer and Ott's argument for the validity of their numerical data (1996) is almost an argument that the basins are r.a. (though they do not use that term). For simpler systems with riddled basins studied previously (Ott, Alexander, et al. 1994), they sketch a procedure to determine, with an arbitrarily small but nonzero chance of error, whether a given point lies in a given basin. The existence of such a pro-

cedure is the central feature of r.a., and in fact the basins in some of those simpler systems can be proven r.a. The authors suppose that the same procedure will work for their new, more realistic example and in fact use it to produce their graphs (Figure 5). It is therefore likely that their basins are r.a., so r.a. does not reflect the undecidability in their dynamical system.

6. The KAM Tori. KAM theory establishes tori of bounded orbits for certain energy-conserving systems. For a given system of the right kind, these tori form “a set of positive measure . . . but a complicated Cantor set” (Moser 1973, 8) akin to the generalized Cantor set that exemplified Theorem IV. Specifically, the union of such tori is nowhere dense, hence riddled, has positive λ -measure, and is therefore u - λ , where λ continues to denote Lebesgue measure (see note 3). This suggests (without proof) that in particular the stability of the solar system may be undecidable in the precise sense that the union of stable orbits is u - λ .

The main KAM theorems concern *Hamiltonian* dynamical systems that are *nearly integrable*. (There are similar theorems for other classes of systems, e.g., Moser 1973, 49.) Hamiltonian systems are those governed by Hamilton's equations,

$$d\mathbf{q}/dt = \partial H/\partial \mathbf{p}, \quad d\mathbf{p}/dt = -\partial H/\partial \mathbf{q}, \quad (2)$$

where H , the *Hamiltonian*, is a function of the $2m$ variables $\mathbf{q} = (q_1, \dots, q_m)$, $\mathbf{p} = (p_1, \dots, p_m)$. For a collection of n point masses, \mathbf{q} might consist of the $3n$ rectangular position coordinates of the bodies and \mathbf{p} the $3n$ momentum components, where H is the total energy of the system; but other coordinates can be chosen.

A Hamiltonian system is *integrable* if \mathbf{q} , \mathbf{p} , and H in (2) can be chosen so that $H = H_0(\mathbf{p})$ is independent of \mathbf{q} . In this case, (2) is easily solved, and all bounded solutions are periodic or *quasi-periodic*, meaning that each is confined to an m -dimensional torus in phase space (which it fills densely), is simple in a certain sense, and will again and again come arbitrarily close to repeating itself (Figure 6).

A *nearly integrable* system is a small perturbation of such a system. For example, a Newtonian system of two gravitating bodies is integrable; adding a very small third body produces a nearly integrable system. Such a perturbation can transform some of the invariant tori into much more complicated sets of chaotic orbits, while other tori are only slightly deformed. KAM theory establishes (among other things) sufficient conditions under which a particular torus is only slightly deformed by a perturbation. The deformed invariant tori so established are called the KAM tori. In the case of an n -body system, the orbits that lie on the KAM tori




Figure 6. Merely schematic illustration of the invariant tori in an integrable system. A quasiperiodic orbit wraps around a torus filling it densely. The cut-away shows such tori filling the phase space.

avoid escape and collision; they are stable in the sense of the historic n -body stability problem.

In order for the methods of KAM theory to show that a particular invariant torus survives a perturbation, the torus must meet certain “nonresonance” conditions.¹¹ Essentially, the partial derivatives $\omega_i = \partial H_0(\mathbf{p})/\partial p_i$ (“frequencies”) of the unperturbed, integrable Hamiltonian $H_0(\mathbf{p})$ on that torus must be far from commensurable, which is usually¹² expressed by a condition of the form

$$\left| \sum_{1 \leq i \leq m} j_i \omega_i \right| \geq 1/\Omega(\sum |j_i|), \quad \forall (j_1, \dots, j_m) \in \mathbf{Z}^m \setminus \{\mathbf{0}\}. \quad (3)$$

If the function Ω is chosen to grow quickly, the set D of frequency vectors $(\omega_1, \dots, \omega_m)$ satisfying this condition has positive λ -measure (see De la Llave 2001). Also, D is nowhere dense, since it is obtained from \mathbf{R}^m by removing a small neighborhood around each m -tuple of commensurable frequencies, and such m -tuples are dense. D is therefore riddled, and by Theorem III, u - λ .

Now, D itself is not the union of the KAM tori in the $2m$ -dimensional phase space. It is a set of m -tuples $(\omega_1, \dots, \omega_m)$ such that each KAM torus *corresponds* to an m -tuple in D . However, this correspondence is such as to ensure that the KAM tori are u - λ . Under mild conditions, there exists a diffeomorphism on an open set $\mathbf{T}^m \times P$ that maps $\mathbf{T}^m \times \Xi \subseteq \mathbf{T}^m \times P$ onto the KAM tori, where \mathbf{T}^m is the m -dimensional torus, P an open

11. Smoothness and nondegeneracy conditions on the Hamiltonian are also required.

12. Arnol'd’s (1963b, 147) nonresonance conditions for the planetary problem are slightly different, but they imply conditions of the form (3) and are implied by other conditions of the form (3), so what is said here applies to them as well.

subset of \mathbf{R}^m , and Ξ a positive-measure subset of D (Pöschel 1982). Since $\mathbf{T}^m \times \Xi$ is nowhere dense and diffeomorphisms preserve this property, the union of KAM tori is nowhere dense. Therefore the latter set is riddled, and the KAM theorems state explicitly that it has positive λ -measure. Hence, the KAM tori themselves form a u - λ set (Figure 7).

In physical applications, the coordinates (\mathbf{q}, \mathbf{p}) in which H_0 is independent of \mathbf{q} will rarely be the most natural phase space coordinates; it is usually necessary to change variables in order to meet the conditions of KAM theorems. However, the coordinates used in KAM theory are always related to the natural coordinates by “canonical” transformations, which preserve both the topology and the natural measure. Hence, when λ is the natural measure, the KAM tori are riddled, positive-measure, and therefore u - λ even in the natural physical coordinates.

Though these tori form u - λ sets, whether the stability problem itself is u - λ is still an open question. There might be many more stable orbits not on the KAM tori, in which case the set of *all* orbits that are stable in our sense might *not* be riddled. However, Arnol’d (1964) has constructed ex-




Figure 7. Schematic cross-section of the invariant tori in a nearly integrable system, showing riddled structure. (In phase spaces with dimension greater than three, orbits are not trapped *inside* such tori.)

amples of orbits of nearly integrable systems that escape, and conjectured that the existence of such orbits is generic. If further *almost* all orbits not lying on the KAM tori escape, then the positive-measure set of stable orbits is riddled and therefore $u\text{-}\lambda$, and the stability of the solar system is undecidable in that concrete sense. Similar claims might also hold for other physical problems subject to KAM theory, such as that of the stability of particle trajectories in an accelerator (Moser 1978).

Incidentally, it is not hard to see that D is r.a., at least if Ω in (3) is Grzegorzczuk-computable and points are coded in a reasonably informative way. The proof idea is just this: To determine with high accuracy whether $(\omega_1, \dots, \omega_m) \in D$, determine whether (3) holds for a large number of integer m -tuples (j_1, \dots, j_m) . Also there should be little difficulty in showing that the KAM tori themselves are r.a. if the relevant transformations are sufficiently computable. Hence again, $u\text{-}\lambda$ characterizes cases of significant undecidability that r.a. obscures.

7. Conclusions. Decidability in a measure μ is a natural relaxation of the concept of decidability, especially in physical contexts. When μ is appropriately chosen, $d\text{-}\mu$ amounts to the existence of a decision procedure that succeeds with probability one. It is also nontrivial in extension and distinct from other relaxed decidabilities.

We have seen that any riddled set with positive Lebesgue measure is undecidable in λ (Theorem III), though such a set may be r.a. (Theorem IV). It therefore appears that $u\text{-}\lambda$ is just the kind of undecidability that Sommerer and Ott intended to assert in 1996. If they are correct in concluding from the numerical evidence that their example has riddled, positive-measure basins, then those basins are $u\text{-}\lambda$, though there is good reason to suspect they are nonetheless r.a.

We have also seen that the KAM tori are $u\text{-}\lambda$, since they too form riddled, positive-measure sets. This suggests that the stability of the solar system, and the qualitative long-term behavior of many other conservative systems, may be undecidable in the concrete sense of $u\text{-}\lambda$. Whether this is so depends on yet unknown facts, but in any case, the concept of undecidability in a measure μ defined here is one rigorous notion of undecidability that can be meaningfully applied to such problems.

One may raise doubts about the physical significance of undecidability results such as those presented here. The models we have considered are highly idealized and grounded in the out-dated prequantum worldview. In particular, questions about the unbounded future of a model may have no physical meaning, since it is intuitively doubtful that any real system will last forever, or more precisely, retain its form—adhere to one model—forever.

There might still be ways in which some actual physical system could

nonetheless behave undecidably in a meaningful sense, but let us not pursue this here. Our primary concern has not been with the occurrence of actual undecidable motion, but with illustrating the concept of decidability in μ . We have not been concerned with the actual solar system, which, due to energy dissipation if nothing else, will no doubt crumble one day. Rather, we have sought to clarify some speculations about a *mathematical* problem raised by classical physics, and to do so in a manner appropriate to that context. More generally, our results on riddled basins and KAM theory have demonstrated fundamental *theoretical* difficulties that arise even for the prediction of simple models, long before real-world contingencies are brought to bear—difficulties quite different from chaos and from quantum indeterminacy. We have seen that *even if* the world were deterministic, classical, susceptible to exact measurement, and well captured by idealized models, some systems could still present significant *computational* barriers to prediction.

Appendix

Let $\|\cdot\|$ denote the Euclidean norm on \mathbf{R}^n . Fix a finite alphabet A and let A^* be the set of finite strings from A . Fix a coding $N: A^* \rightarrow \mathbf{N}$ onto \mathbf{N} , and for each n , a coding $Q_n: A^* \rightarrow \mathbf{Q}^n$ onto \mathbf{Q}^n , such that the relation $\|Q_n(s_1) - Q_n(s_2)\| \leq Q_n(s_3)$ is recursive.

Definition 1.

- (i) A *Cauchy oracle* for x is a function $\phi: A^* \rightarrow A^*$ such that $\|Q_n(\phi(s)) - x\| < 2^{-N(s)}$ for all $s \in A^*$. CO_x denotes the set of Cauchy oracles for x .
- (ii) When an OTM M is supplied with a particular oracle ϕ we refer to it as M^ϕ . If M^ϕ halts on an input string s , we write $M^\phi(s)\downarrow$ and denote the output string by $M^\phi(s)$ or $M^\phi(s)\downarrow$.
- (iii) Let $\sigma: \{0, 1, \dots, k\} \rightarrow A^*$ be a finite sequence of strings. We say that $\text{length}(\sigma) = k + 1$. For any $\phi: \mathbf{N} \rightarrow A^*$, we write $\sigma \subset \phi$ and $\phi \supset \sigma$ if $\sigma(i) = \phi(i)$ for all $i \in \text{dom}(\sigma) = \{0, 1, \dots, k\}$.
- (iv) We write $M^\sigma(n)\downarrow = x$ if for some oracle function $\phi \supset \sigma$, $M^\phi(n)\downarrow = x$ and $M^\phi(n)$ queries the oracle only with numbers in $\text{dom}(\sigma)$ —i.e., M^ϕ never enters the query state with a string s on the oracle tape such that $N(s) \geq \text{length}(\sigma)$. If no such ϕ exists, we write $M^\sigma(n)\uparrow$.

Proposition 1 (Use Principle). $M^\phi(n)\downarrow = x \Leftrightarrow (\exists \sigma \subset \phi) M^\sigma(n)\downarrow = x$.

Proof. See Soare (1987). ■

Definition 2. $B(x, \varepsilon)$ ($B[x, \varepsilon]$) denotes the open (closed) ball with center x and radius ε .

Theorem I (Topological Use Principle). If $M^\phi(n)\downarrow = q$ for some $\phi \in CO_x$, then there is a neighborhood U of x such that $(\forall y \in U)(\exists \psi \in CO_y) M^\psi(n)\downarrow = q$.

Proof. Fix $x \in \mathbf{R}^n$ and suppose ϕ satisfies the antecedent. By the Use Principle, choose a finite sequence σ such that $\sigma \subset \phi$ and $M^\sigma(n)\downarrow = q$. Let $U = \bigcap_{i \leq \text{length}(\sigma)} B(\phi(i), 2^{-i})$. To see that U satisfies the consequent, let $y \in U$. Choose $\theta \in CO_y$ and let

$$\psi(i) = \begin{cases} \sigma(i) & \text{if } i \leq \text{length}(\sigma) \\ \theta(i) & \text{if } i > \text{length}(\sigma). \end{cases}$$

Then $\psi \in CO_y$, and by the Use Principle, $M^\psi(n)\downarrow = M^\sigma(n)\downarrow = q$. ■

Definition 3. A set $B \subseteq \mathbf{R}^n$ is *recursively approximable (r.a.)* if there is an OTM M such that $(\forall x \in \mathbf{R}^n, \phi \in CO_x, m \in \mathbf{Z}^+)(M^\phi(m)\downarrow \text{ and } \lambda^*E_{M,m}(B) \leq 2^{-m})$, where λ^* denotes Lebesgue outer measure and $E_{M,m}(B) = \{x \in \mathbf{R}^n \mid (\exists \phi \in CO_x) M^\phi(m) \neq \chi_B(x)\}$.

Definition 4. For any measure μ on \mathbf{R}^n , a set $B \subseteq \mathbf{R}^n$ is *decidable in μ* (or $d\text{-}\mu$) if there exists an OTM M such that

$$E_M(B) = \left\{ x \in \mathbf{R}^n \mid (\exists \phi \in CO_x) [M^\phi(\emptyset)\downarrow \neq \chi_B(x) \text{ or } M^\phi(\emptyset)\uparrow] \right\}$$

is μ -measurable and $\mu E_M(B) = 0$. (The null string \emptyset here is an arbitrary choice.) Otherwise, B is *undecidable in μ* (or $u\text{-}\mu$).

Note that without the alternative $M^\phi(\emptyset)\uparrow$ here, only measure-theoretically trivial sets would be $d\text{-}\mu$:

Proposition 2. Suppose μ^* is a regular outer measure on \mathbf{R}^n (i.e., if $U \neq \emptyset$ is open in the Euclidean topology then $\mu^*U > 0$), $B \subseteq \mathbf{R}^n$, M is an OTM, $\mu E_M(B) = 0$, and $(\forall x \in \mathbf{R}^n)(\forall \phi \in CO_x) M^\phi(\emptyset)\downarrow \in \{0, 1\}$. Then $\mu B = 0$ or $\mu(B^c) = 0$, where $B^c = \mathbf{R}^n \setminus B$.

Proof. Assume antecedents. Let $U = \{x \mid (\exists \phi \in CO_x) M^\phi(\emptyset)\downarrow = 1\}$, $V = \{x \mid (\exists \phi \in CO_x) M^\phi(\emptyset)\downarrow = 0\}$. By Theorem I, U and V are open. Since $(\forall x \in \mathbf{R}^n)(\forall \phi \in CO_x) M^\phi(\emptyset)\downarrow \in \{0, 1\}$, $U \cup V = \mathbf{R}^n$. Since \mathbf{R}^n is connected, either $U = \emptyset$, $V = \emptyset$, or $U \cap V \neq \emptyset$. But $U \cap V \subseteq E_M(B)$, so $\mu(U \cap V) = 0$. Since $U \cap V$ is open and μ^* is regular, $U \cap V = \emptyset$. Therefore $U = \emptyset$ or $V = \emptyset$. Hence $B \subseteq E_M(B)$ or $B^c \subseteq E_M(B)$. Since $\mu E_M(B) = 0$, either $\mu B = 0$ or $\mu(B^c) = 0$. ■

Theorem II. If $B \subseteq \mathbf{R}^n$ is d- λ then B is r.a.

Proof. See main text for strategy. Let $S: \mathbf{N} \rightarrow A^{**}$ be an effective enumeration of the finite sequences of strings in A^* . Define $U(i)$ as the set of points with Cauchy oracles ϕ such that $S(i) \subset \phi$, i.e., $U(i) = \bigcap_{j \leq \text{length}(S(i))} B(Q_n(S(i)_j), 2^{-j})$ where $S(i)_j$ is the j^{th} string in $S(i)$. Where $\text{length}(S(i)) = 2j$, let $V(i) = \bigcup_{k \leq j} B(Q_n(S(i)_{2k-1}), Q_n(S(i)_{2k}))$.

Suppose an OTM M satisfies Definition 4. For each $m \in \mathbf{N}$, construct recursive sequences of integers $u[m, i], v[m, i]$ as follows:

1. Set $R, i = 1$.
2. Simulate $M^{S(i)}$ for all $j \in \mathbf{N}$ simultaneously (by dovetailing). Whenever $M^{S(i)}(\emptyset) \downarrow$ for some j , let $u[m, i] = j$ and $i = i + 1$. Once $\lambda(B[0, R] \setminus \bigcup_{k \leq i} U(u[m, k])) < 2^{-m-R-1}$, proceed to 3. (Volumes of finite unions, intersections, and differences of rational balls can be computed to any accuracy, since we assume $\|Q_n(s_1) - Q_n(s_2)\| \leq Q_n(s_3)$ is recursive.)
3. By trial and error, fix $v[m, R]$ such that $\lambda V(v[m, R]) < 2^{-m-R-1}$ and $B[0, R] \subseteq \bigcup_{j \leq i} U(u[m, j]) \cup \bigcup_{j \leq R} V(v[m, j])$. Let $R = R + 1$ and go to 2.

Each repetition of 2 and 3 will halt because $B[0, R]$ is compact. Note $\bigcup_i [U(u[m, i]) \cup V(v[m, i])] = \mathbf{R}^n$, and $\lambda \bigcup_i V(v[m, i]) < 2^{-m}$.

Now an algorithm for a machine M' satisfying Definition 3 proceeds thus: given input m and oracle $\phi \in CO_x$, evaluate for each i (again by dovetailing) whether $x \in U(u[m, i])$ and whether $x \in V(v[m, i])$. If $x \in U(u[m, i])$, output $M^{S(u[m, i])}(\emptyset)$. If $x \in V(v[m, i])$, output 0.

Since $\bigcup_i [U(u[m, i]) \cup V(v[m, i])] = \mathbf{R}^n$, M' will halt on every Cauchy oracle. Also, $E_{M', m}(B) \subseteq E_M(B) \cup \bigcup_i V(v[m, i])$. Since $\lambda E_M(B) = 0$ and $\lambda \bigcup_i V(v[m, i]) < 2^{-m}$, $\lambda E_{M', m}(B) < 2^{-m}$. ■

Definition 5. A set $B \subseteq \mathbf{R}^n$ is *riddled* if for every open set $U \neq \emptyset$ in the Euclidean topology, $\lambda^*(U \setminus B) > 0$.

Theorem III. If $B \subseteq \mathbf{R}^n$ is riddled and $\lambda^*B > 0$, then B is u- λ .

Proof. Assume the antecedents and suppose M is an OTM. We show that $\lambda^*E_M(B) > 0$.

Case 1: $\neg(\exists x \in \mathbf{R}^n)(\forall \phi \in CO_x) M^\phi(\emptyset) = 1$. Then $E_M(B) \supseteq B$, so $\lambda^*E_M(B) \geq \lambda^*B > 0$.

Case 2: $(\exists x \in \mathbf{R}^n)(\forall \phi \in CO_x) M^\phi(\emptyset) = 1$. Then by Theorem I, there is a neighborhood U of x such that $(\forall y \in U)(\exists \psi \in CO_y) M^\psi(\emptyset) \downarrow = 1$. Therefore $U \setminus B \subseteq E_M(B)$. But by riddling, $\lambda^*(U \setminus B) > 0$. Therefore $\lambda^*E_M(B) > 0$, so B is u- λ . ■

Theorem IV. There exists a subset of \mathbf{R} that is r.a. but not d- λ .

Proof. Say a closed interval I is *maximal* in a set $S \subseteq \mathbf{R}$ if $I \subseteq S$ and for every closed interval $J \subseteq S, J \cap I \neq \emptyset \Rightarrow J \subseteq I$. We construct a generalized Cantor set C as follows:

- (i) Let $C_0 = [0, 1]$.
- (ii) For each $i \in \mathbf{Z}^+$, let $C_i = C_{i-1} \setminus \cup \{B([a + b]/2, 2^{-i} \cdot [b - a]) \mid [a, b] \text{ is maximal in } C_{i-1}\}$.
- (iii) Let $C = \cap_i C_i$.

Part (ii) dictates that we obtain C_i by removing the middle $(2^i)^{\text{th}}$ of each maximal interval in C_{i-1} . The set C is the limit of this process, and is clearly riddled by construction.

We show that $\lambda C > 0$. Since the C_i are all measurable, $\lambda C = 1 - \sum_i \lambda(C_{i-1} \setminus C_i)$. Our construction (ii) removes the middle $(2^i)^{\text{th}}$ part from each component of C_{i-1} . Therefore, $\lambda(C_{i-1} \setminus C_i) = 2^{-i} \lambda C_{i-1} \leq 2^{-i}$, with strict inequality for $i > 1$. So $\sum_i \lambda(C_{i-1} \setminus C_i) < \sum_i 2^{-i} = 1$. Thus $\lambda C > 0$. By Theorem III, C is u- λ .

We now show that C is r.a. by sketching an OTM that approximates C . The algorithm is this: given input m , let M^ϕ request from its oracle ϕ the value of $\phi(2m + 3)$, and determine whether the $Q[\phi(2m + 3)]$ is in C_{m+1} . (This can be done effectively since one can construct the rational endpoints of C_{m+1} following (i)-(iii), and we code rationals so that $|q_1 - q_2| \leq q_3$ is recursive.) If $Q[\phi(2m + 3)] \in C_{m+1}$, let $M^\phi(m) = 1$; otherwise $M^\phi(m) = 0$.

Since $|Q[\phi(2m + 3)] - x| < 2^{-(2m+3)}$ any $x \in E_{M, m}(C_{m+1})$ must be close to one of the 2^{m+2} endpoints of C_{m+1} , within distance $2^{-(2m+3)}$. Therefore $\lambda^* E_{M, m}(C_{m+1}) \leq (2^{m+2})(2^{-(2m+3)}) = 2^{-(m+1)}$. So, if $\lambda(C_{m+1} \setminus C) \leq 2^{-(m+1)}$, then $\lambda^* E_{M, m}(C) \leq 2^{-(m+1)} + 2^{-(m+1)} = 2^{-m}$. This is in fact the case, for we now show that for all $i, \lambda(C_i \setminus C) < 2^{-i}$. For $i = 0$ we have $\lambda(C_0 \setminus C) = \lambda C_0 - \lambda C < 1 = 2^0$, since $\lambda C > 0$. For $i > 0$ we have $\lambda(C_i \setminus C) = \sum_{j>i} \lambda(C_{j-1} - C_j) =$ (by construction) $\sum_{j>i} 2^{-j} \lambda C_{j-1} < \sum_{j>i} 2^{-j} = 2^{-i}$. Therefore $\lambda^* E_{M, m}(C) \leq 2^{-m}$, so C is r.a. ■

REFERENCES

Alexander, J. C., James A. Yorke, Zhiping You, and Ittai Kan (1992), "Riddled Basins", *International Journal of Bifurcation and Chaos* 2: 795–813.
 Arnol'd, Vladimir I. (1963a), "Proof of a Theorem of A. N. Kolmogorov on the Invariance of Quasi-Periodic Motions Under Small Perturbations of the Hamiltonian", *Russian Mathematical Surveys* 18: 9–36.
 ——— (1963b), "Small Denominators and Problems of Stability of Motion in Classical and Celestial Mechanics", *Russian Mathematical Surveys* 18: 85–191.
 ——— (1964), "Instability of Dynamical Systems with Several Degrees of Freedom", *Soviet Mathematics* 5: 581–585.

- Batterman, Robert W. (1998), "Why Equilibrium Statistical Mechanics Works: Universality and the Renormalization Group", *Philosophy of Science* 65: 183–208
- Blum, Lenore, Mike Shub, and Steve Smale (1989), "On a Theory of Computation and Complexity over the Real Numbers: NP-Completeness, Recursive Functions, and Universal Machines", *Bull. Am. Math. Soc.* 21: 1–46.
- Davis, Martin D., and Elaine J. Weyuker (1983), *Computability, Complexity, and Languages*. San Diego, CA: Academic Press.
- De la Llave, Rafael (2001), "A Tutorial on KAM Theory", *Mathematical Physics Archive* (01–29), http://www.ma.utexas.edu/mp_arc/index-01.html.
- Diacu, Florin, and Philip Holmes (1996), *Celestial Encounters: The Origins of Chaos and Stability*. Princeton: Princeton University Press.
- Gödel, Kurt (1931), "Über Formal Unentscheidbare Studer *Principia Mathematica und verwandter Systeme I*", *Monatshefte fur Mathematik und Physik* 38: 173–198. Translated in Kurt Gödel (1986), *Collected Works I*. Oxford: Oxford University Press.
- Goroff, Daniel L. (1993), Introduction to *New Methods of Celestial Mechanics* by Henri Poincaré. Woodbury, NY: American Institute of Physics.
- Grzegorzczak, A. (1955a), "Computable Functionals", *Fundamenta Mathematicae* 42: 169–202.
- (1955b), "On the Definition of Computable Functionals", *Fundamenta Mathematicae* 42: 233–239.
- (1957), "On the Definitions of Computable Real Continuous Functions", *Fundamenta Mathematicae* 44: 61–67.
- Heagy, James F., Thomas L. Carroll, and Louis M. Pecora (1994), "Experimental and Numerical Evidence for Riddled Basins in Coupled Chaotic Systems", *Physical Review Letters* 73: 3528–3531.
- Ko, Ker-I (1991), *Complexity Theory of Real Functions*. Boston: Birkhauser.
- Malament, David, and Sandy Zabell (1980), "Why Gibbs Phase Averages Work—The Role of Ergodic Theory", *Philosophy of Science* 47: 339–349.
- Milnor, John (1985), "On the Concept of Attractor", *Communications in Mathematical Physics* 99: 177–195.
- Moore, Christopher (1990), "Unpredictability and Undecidability in Dynamical Systems", *Physical Review Letters* 64: 2354–2357.
- (1991), "Generalized Shifts: Unpredictability and Undecidability in Dynamical Systems", *Nonlinearity* 4: 199–230.
- Moser, Jürgen (1973), *Stable and Random Motions in Dynamical Systems*. Princeton: Princeton University Press.
- (1978), "Is the Solar System Stable?", *Mathematical Intelligencer* 1: 65–71.
- Myrvold, Wayne C. (1997), "The Decision Problem for Entanglement," in R. S. Cohen et al. (eds.), *Potentiality, Entanglement, and Passion-at-a-Distance*. Great Britain: Kluwer Academic Publishers, 177–190.
- Ott, Edward, J. C. Alexander, Ittai Kan, John C. Sommerer, and James A. Yorke (1994), "The Transition to Chaotic Attractors with Riddled Basins", *Physica D* 76: 384.
- Parker, Matthew W. (1998), "Did Poincaré Really Discover Chaos?", *Stud. Hist. Phil. Mod. Phys.* 29, 578–588.
- Pitowsky, Itamar (1996), "Laplace's Demon Consults an Oracle: The Computational Complexity of Prediction", *Stud. Hist. Phil. Mod. Phys.* 27: 161–180.
- Poincaré, Henri (1890), "Sur le Problème des Trois Corps et les Équations de la Dynamique", *Acta Mathematica* 13: 1–270.
- (1891), "Le Problème des Trois Corps", *Revue Générale des Sciences Pures et Appliquées* 2 (1): 1–5.
- (1898), "On the Stability of the Solar System", *Nature* 58: 183.
- (1907) "Le Hasard", *La Revue du Mois* 3: 257–276. Translated in Henri Poincaré (1952), *Science and Method*. New York: Dover Publications, Inc.
- ([1892–1899] 1993) *New Methods of Celestial Mechanics*. Reprint. Woodbury, NY: American Institute of Physics. Originally published as *Les Méthodes Nouvelles de la Mécanique Céleste*.
- Pöschel, Jürgen (1982) "Integrability of Hamiltonian Systems on Cantor Sets", *Communications on Pure and Applied Mathematics* 35: 653–695.

- Pour-El, Marian Boykan, and Jerome Caldwell (1975), "On a Simple Definition of Computable Function of a Real Variable—With Application to Functions of a Complex Variable", *Zeitschr. f. math. Logik und Grundlagen d. Math.* 21: 1–19.
- Pour-El, Marian Boykan, and Ian Richards (1983), "Computability and Noncomputability in Classical Analysis," *Trans. Am. Math. Soc.* 275: 539–545.
- Sklar, Lawrence (1973), "Statistical Explanation and Ergodic Theory", *Philosophy of Science* 40: 194–212.
- (1993), *Physics and Chance*. Cambridge: Cambridge University Press.
- Soare, Robert I. (1987), *Recursively Enumerable Sets and Degrees*. New York: Springer-Verlag.
- Sommerer, John C., and Edward Ott (1993), "A Physical System with Qualitatively Uncertain Dynamics", *Nature* 365: 138–140.
- (1996), "Intermingled Basins of Attraction: Uncomputability in a Simple Physical System", *Physics Letters A* 214: 243–251. Figure reprinted with permission from Elsevier Science, Copyright 1996.
- Turing, Alan (1936), "On Computable Numbers with an Application to the Entscheidungsproblem", *Proc. London Math. Soc* 42: 230–265.
- Vranas, Peter B. M. (1998), "Epsilon-Ergodicity and the Success of Equilibrium Statistical Mechanics", *Philosophy of Science* 65: 688–708.
- Weihrauch, Klaus (2000), *Computable Analysis*. Berlin: Springer-Verlag.
- Wolfram, Stephen (1985), "Undecidability and Intractability in Theoretical Physics", *Physical Review Letters* 54: 735–738.